

Biological Control Strategies in Integrated Pest Management

Mobashshir Kabiraj^{*1}, Anisha Parvin², Karan Kashyap³ and Ansh Sharma⁴

¹M.Sc. (Ag.) Entomology, Division of Entomology, Faculty of Agriculture, Sher-e-Kashmir University of Agricultural Sciences and Technology of Kashmir

²M.Sc. (Agri.) Plant Pathology, Department of Plant Pathology, School of Agriculture, Lovely Professional University, Phagwara, Punjab

³M.Sc. (Agri.) Entomology, Department of Entomology, School of Agriculture, Lovely Professional University, Phagwara, Punjab

⁴M.Sc. (Agri.) Entomology, Department of Entomology, School of Agriculture, Lovely Professional University, Phagwara, Punjab

*Corresponding Author Email: mobashshirkabiraj124@gmail.com

Abstract

Biological control is a core ecological component of integrated pest management (IPM) because it uses predators, parasitoids, microbial agents and other natural enemies to suppress pest populations while reducing unnecessary dependence on broad-spectrum pesticides. Its value lies not only in replacing a chemical treatment, but in strengthening the biological regulation that operates within crop ecosystems throughout the season. Conservation of resident natural enemies, planned releases of mass-reared predators or parasitoids, microbial insecticides and antagonistic microorganisms can all contribute to pest suppression when they are matched with the target pest, crop stage and local environment. Biological control is most effective when supported by regular scouting, economic thresholds, correct timing, habitat management, resistant varieties, sanitation and selective pesticide use. Important agents include ladybird beetles, lacewings, predatory mites, Trichogramma parasitoids, *Bacillus thuringiensis*, entomopathogenic fungi and baculoviruses. However, biological control is not automatically successful. Temperature, humidity, pesticide residues, product quality, pest stage and release timing can strongly influence performance, and introductions intended for permanent establishment require careful ecological risk assessment. New developments in molecular identification, mass rearing, microbial formulation, digital monitoring and precision release are making biological control more reliable and easier to integrate with other tactics.

Keywords: Biological control, Integrated pest management, Natural enemies, Parasitoids, Predators, Microbial agents, Sustainable agriculture

How to Cite this article: Kabiraj, M., Parvin, A., Kashyap, K., & Sharma, A. (2026). Biological Control Strategies in Integrated Pest Management. *Future Agriculture e-magazine*, September 2026, Issue 3, pp. 198-203.

Introduction

Integrated pest management (IPM) is built on the idea that crop protection should not depend on a single method. Instead, farmers combine prevention, monitoring, cultural practices, resistant varieties, mechanical measures, biological control and carefully selected pesticides so that pest populations remain below damaging levels. Biological

control is one of the most important ecological components of this approach because it uses living organisms or their products to suppress

pests. Predators, parasitoids, pathogenic microorganisms and naturally occurring antagonists can reduce pest pressure while helping to conserve the wider farm ecosystem.

The value of biological control becomes especially clear when pest management is viewed over an entire season rather than as a

single spray decision. Broad-spectrum pesticide use may provide rapid knockdown, but repeated use can disrupt beneficial organisms, select for resistance and allow secondary pests to become more important. Biological control works differently. It aims to strengthen the regulating forces that are already present in agricultural ecosystems or to introduce carefully selected beneficial organisms where natural regulation is inadequate.

A successful IPM programme therefore asks two questions at the same time: how many pests are present, and what is happening to their natural enemies? Decisions based only on pest counts can miss an important part of the field situation. A moderate pest population may decline naturally when predators or parasitoids are abundant, while a smaller population may increase rapidly if beneficial organisms have been destroyed. Biological control makes IPM more responsive to these ecological relationships.

Biological Control within the IPM

Biological control can be broadly described as the use of natural enemies to reduce the population, activity or damage caused by a pest. In IPM it is rarely used in isolation. Instead, it is integrated with agronomic practices that make the crop less favourable to pests and more favourable to beneficial organisms. Crop rotation, sanitation, balanced fertilization, appropriate planting time, trap crops, habitat management and selective pesticide use can all influence the success of natural enemies.



The objective is not necessarily to eliminate every pest individual. Complete eradication is usually unnecessary and may even be ecologically unrealistic. IPM focuses on

keeping pest numbers below the economic threshold, the level at which action is justified to prevent unacceptable loss. Biological control contributes by slowing pest population growth, reducing peak densities and lowering the frequency with which chemical intervention is required.

This preventive role is important. Natural enemies often work continuously in the background and may suppress pests before crop damage becomes obvious. Their effect can be difficult to notice because the result is the absence of an outbreak. For this reason, scouting should record predators, parasitoid activity, parasitized eggs or larvae, diseased insects and other signs of natural regulation, not simply count the target pest.

Predators as Natural Pest Regulators

Predators consume several prey individuals during their life and can act across different pest stages. Ladybird beetles, lacewings, predatory bugs, spiders, mites and many other groups are common examples in agricultural systems. Their importance varies with crop, climate and pest complex, but they often contribute to suppression of aphids, whiteflies, thrips, mites, caterpillars, eggs and other soft-bodied pests.

Predator conservation begins with reducing avoidable mortality. Unnecessary broad-spectrum sprays can remove beneficial organisms faster than pests recover, particularly when pest species reproduce rapidly or migrate into the field. Selective products, spot treatment, need-based application and correct timing can help preserve predator populations. Flowering borders, refuges and diversified vegetation may provide pollen, nectar, shelter or alternative prey, but habitat manipulation must be planned carefully so that it does not also favour key pests.

Predators are especially useful as part of a community of natural enemies. One species may feed mainly on eggs, another on early larvae and another on mobile adults. This complementary action can make biological regulation more stable. However, predator abundance alone should not be treated as a guarantee of control. Field observations are still needed to determine whether predation is

sufficient to keep the pest below the action threshold.

Parasitoids and Their Role in Targeted Suppression

Parasitoids are insects whose immature stages develop on or inside a single host and ultimately kill it. Many parasitoid wasps and flies are highly specialized, making them valuable against particular crop pests. Egg parasitoids such as *Trichogramma* species are widely recognized in crop protection because they attack the pest before the damaging larval stage emerges. Other parasitoids attack larvae, pupae or adults.


Parasitoid performance depends strongly on timing. Releasing an egg parasitoid after most eggs have already hatched gives little benefit. Monitoring pest phenology, moth activity, egg laying and crop stage therefore improves release decisions. In augmentative biological control, beneficial insects may be mass-produced and released in inoculative or inundative programmes. Release rate, interval, distribution across the field and product quality all affect results.

Field conditions also matter. Extreme temperature, heavy rain, pesticide residues and a shortage of food resources for adult parasitoids can reduce survival. Conserving flowering plants that provide accessible nectar can support some species, while selective pesticide programmes can prevent unnecessary mortality. Compatibility between biological releases and other IPM tactics should be considered before the season begins rather than after problems occur.

Microbial Biological Control Agents

Microorganisms provide another major group of biological control tools. Bacteria, fungi, viruses and other microbes can infect or antagonize pests. *Bacillus thuringiensis* is one of the best-known microbial insecticides; its protein toxins are active against susceptible insect groups when used correctly. Entomopathogenic fungi such as *Beauveria bassiana* and *Metarhizium* species infect insects through the cuticle, while nucleopolyhedroviruses can be highly specific to particular caterpillar pests.

Microbial agents can fit well into IPM because many are selective and leave predators and parasitoids relatively undisturbed compared with broad-spectrum insecticides. Their performance, however, is closely linked to environmental conditions. Ultraviolet light, temperature, humidity, rainfall, formulation, spray coverage and the developmental stage of the target pest can all influence field efficacy. For example, many microbial products perform best when applied against young pest stages before severe crop damage has occurred.



Biological control is also important in plant disease management. Antagonistic fungi and bacteria can compete with pathogens, produce inhibitory substances, parasitize pathogen structures or stimulate plant defence responses. Products based on *Trichoderma* and beneficial bacteria are used in several crop systems as seed, soil or root-zone treatments. As with insect biocontrol, quality, viable propagule count, storage and correct application are essential for dependable results.

Classical, Augmentative and Conservation Biological Control

Biological control strategies are often grouped into classical, augmentative and conservation approaches. Classical biological control involves the introduction and establishment of a natural enemy against an invasive or introduced pest, normally after detailed study of host specificity and environmental risk. When successful, the natural enemy can provide long-term suppression, but such programmes require rigorous scientific and regulatory assessment.

because the organism is intended to establish in the environment.

Augmentative biological control uses repeated releases of natural enemies that are already known to be useful. Inoculative releases introduce relatively small numbers early so that the beneficial population can increase, whereas inundative releases use larger numbers for more immediate suppression. Commercial production of parasitoids, predators and microbial agents has made this approach increasingly practical in protected cultivation and in some field crops.

Conservation biological control focuses on protecting and strengthening beneficial organisms already present in the agroecosystem. It may include reducing disruptive pesticide use, providing shelter or floral resources, maintaining field margins and improving crop diversity. Conservation is often the most economical starting point because the natural enemies are already adapted to local conditions. The goal is to avoid destroying an ecosystem service that is available at no additional purchase cost.

Monitoring, Thresholds and Decision Making

Biological control works best when supported by regular field monitoring. Scouting should be systematic and should record pest density, crop damage, beneficial organisms and environmental conditions. Sticky traps, pheromone traps, light traps, plant inspections and direct counts can help identify changes in pest activity. The information is then compared with action thresholds or locally validated decision rules.

Timing is critical because natural enemies often need time to act. A farmer who waits until the crop is heavily damaged may conclude that biological control is too slow, when the real problem is delayed intervention. Early detection allows parasitoids, predators or microbial agents to be used against more susceptible pest stages. It also allows chemical measures, when necessary, to be chosen in ways that preserve biological agents.

Digital scouting tools, mobile applications, remote sensing and image-based identification are beginning to improve monitoring. Their role

is not to replace field knowledge but to make records more consistent and help identify trends. When pest and natural-enemy data are stored over several seasons, farms can better understand recurring hotspots and the conditions under which biological control succeeds or fails.

Compatibility with Pesticides and Other IPM Tactics

Biological and chemical control do not have to be opposing strategies. In well-designed IPM they can be compatible when pesticide selection and timing are based on need. Selective insecticides, biopesticides, insect growth regulators and localized treatments may preserve more beneficial organisms than routine broad-spectrum applications. The pesticide label, known effects on natural enemies, residual activity and crop stage should be considered before treatment.

Cultural tactics can also strengthen biological control. Removal of infested crop residues can reduce carry-over populations, while crop rotation interrupts pest life cycles. Balanced nitrogen management is important because excessive vegetative growth can sometimes favour sap-feeding pests. Resistant varieties reduce pest establishment or damage and can give natural enemies more time to act. Physical barriers, traps and sanitation further reduce the population that biological agents must suppress.

The strongest IPM programmes use several moderate pressures rather than expecting one tactic to solve the whole problem. A pest population weakened by host-plant resistance, cultural practices and natural enemies may require fewer pesticide interventions. This reduces selection pressure for resistance and helps maintain the effectiveness of both chemical and biological tools.

Table 1. Major biological control approaches used in integrated pest management.

Approach	Role in IPM
Conservation biological control	Protects resident predators, parasitoids and antagonists through selective inputs and supportive habitat.
Augmentative release	Adds mass-reared natural enemies at critical crop stages to reinforce pest suppression.
Microbial control	Uses organisms such as <i>Bacillus thuringiensis</i> , <i>Beauveria</i> , <i>Metarhizium</i> or baculoviruses against susceptible pests.

Approach	Role in IPM
Classical biological control	Introduces carefully selected natural enemies for long-term suppression of invasive pests after risk assessment.
Biological disease suppression	Uses beneficial fungi or bacteria to reduce pathogen survival, infection or root-zone disease pressure.

Advantages of Biological Control in Sustainable Agriculture

Biological control can provide important environmental and economic benefits. By reducing reliance on repeated broad-spectrum pesticide use, it can help conserve pollinators and other non-target organisms, reduce pesticide residues and slow the development of resistance. Established natural enemies may continue to provide suppression across seasons, while microbial and augmentative products offer options where residue restrictions or worker-safety considerations are important.

The approach also fits the wider goal of ecological intensification, in which biological processes are used to support production. Healthy predator and parasitoid communities contribute to resilience because they can respond when pests begin to increase. This does not mean that biological control is free or effortless. Monitoring, identification, habitat management and quality biological products require knowledge and investment. The benefit is a more balanced system with fewer unnecessary interventions.

For farmers, the economic value should be assessed at the programme level. A biological product may appear more expensive than a single conventional spray, but the comparison changes if it reduces later treatments, prevents resistance problems, protects beneficial organisms or supports market requirements for lower residues. Local cost, crop value and expected pest pressure must therefore be included in decision making.

Table 2. Compatibility of biological control with other IPM components.

IPM component	Compatibility point
Monitoring and thresholds	Use scouting, traps and natural-enemy counts so intervention is timed only when needed.
Selective pesticides	Choose products and timings with lower impact on beneficial organisms; avoid disruptive broad-spectrum residues.
Habitat management	Provide flowering resources or refuges where appropriate while avoiding pest or disease reservoirs.

IPM component	Compatibility point
Cultural control	Coordinate sanitation, rotation and resistant cultivars so they lower pest pressure without unnecessarily destroying beneficials.
Behavioural tools	Pheromone monitoring, mating disruption or attract-and-kill can complement releases and microbial agents.

Limitations and Practical Challenges

Biological control is not equally reliable under all conditions. Natural enemies are living organisms and their activity changes with temperature, humidity, crop structure, pesticide exposure and availability of hosts or prey. Some agents act more slowly than chemical insecticides, which can be a disadvantage when pest density is already far above the economic threshold. Biological products also require careful storage and handling to maintain viability.

Correct identification is another challenge. Releasing the wrong natural enemy or applying a microbial product against a non-susceptible pest stage can waste time and money. Product quality is equally important. Commercial cultures must have adequate survival, vigour and biological activity, and field releases should be distributed uniformly. Farmers and advisers therefore need reliable suppliers and practical guidance.

Ecological risk must also be considered, especially for introductions intended to establish permanently. Modern biological control programmes emphasize host-range testing, regulatory review and post-release monitoring. The objective is to obtain pest suppression without unacceptable effects on non-target species. Careful selection and responsible use are essential parts of biological control, not obstacles to it.

Emerging Directions in Biological Control

New technologies are making biological control more precise. Molecular tools can improve identification of pests, parasitoids and microbial strains, while population genetics can help trace establishment and dispersal. Improved mass-rearing systems are increasing the consistency of commercial natural enemies. Formulation research is helping microbial agents survive storage, ultraviolet exposure and field application.

Semiochemicals and behavioural tools can also be combined with biological control. Pheromone traps may indicate the best release time for parasitoids, and attract-and-retain approaches may help concentrate beneficial insects where they are needed. In protected cultivation, banker plants and managed flowering resources can sustain natural enemies between pest outbreaks. Automated release devices and drones are being explored for distributing some biological agents over larger areas.

The future of biological control within IPM will depend less on a single new organism and more on better system design. Reliable monitoring, quality-controlled agents, selective pesticides, habitat management, farmer training and locally validated thresholds need to work together. Biological control is strongest when it is treated as a central management process rather than an emergency substitute after other methods fail.

Conclusion

Biological control is a practical and scientifically grounded component of integrated pest management. Predators, parasitoids, microbial agents and conserved natural enemies can suppress pests while reducing dependence on repeated broad-spectrum pesticide use. Their success depends on correct identification, early monitoring, appropriate timing, good product quality and compatibility with cultural, physical and chemical tactics.

The most effective strategy is not to replace every pesticide with a biological product, but to design a crop-protection system in which biological regulation is protected and used whenever it can contribute. Conservation of beneficial organisms, threshold-based decisions and selective interventions make IPM more durable. As monitoring, mass production, microbial formulation and digital decision tools improve, biological control is likely to become even more important in productive, resilient and environmentally responsible agriculture.

References

Bale, J. S., van Lenteren, J. C., & Bigler, F. (2008). Biological control and sustainable food production. *Philosophical Transactions of the*

Royal Society B: Biological Sciences, 363(1492), 761-776.

Deguine, J. P., Aubertot, J. N., Flor, R. J., Lescouret, F., Wyckhuys, K. A. G., & Ratnadass, A. (2021). Integrated pest management: Good intentions, hard realities. A review. *Agronomy for Sustainable Development*, 41, 38.

Eilenberg, J., Hajek, A., & Lomer, C. (2001). Suggestions for unifying the terminology in biological control. *BioControl*, 46, 387-400.

Hajek, A. E., & Eilenberg, J. (2018). *Natural Enemies: An Introduction to Biological Control* (2nd ed.). Cambridge University Press.

Heimpel, G. E., & Mills, N. J. (2017). *Biological Control: Ecology and Applications*. Cambridge University Press.

Lacey, L. A., Grzywacz, D., Shapiro-Ilan, D. I., Frutos, R., Brownbridge, M., & Goettel, M. S. (2015). Insect pathogens as biological control agents: Back to the future. *Journal of Invertebrate Pathology*, 132, 1-41.

Pretty, J., & Bharucha, Z. P. (2015). Integrated pest management for sustainable intensification of agriculture in Asia and Africa. *Insects*, 6(1), 152-182.

Stenberg, J. A. (2017). A conceptual framework for integrated pest management. *Trends in Plant Science*, 22(9), 759-769.

van Lenteren, J. C., Bolckmans, K., Köhl, J., Ravensberg, W. J., & Urbaneja, A. (2018). Biological control using invertebrates and microorganisms: Plenty of new opportunities. *BioControl*, 63, 39-59.