

Advanced Diagnostic Technologies in Veterinary Practice

S.S. Mehesare^{*1} and S. M. Khupse²

¹Assistant Professor, Department of Veterinary Clinical Complex (Veterinary Medicine), Mahatma Gandhi Mission, MGM's Dr. Rajendra Suryawanshi College of Veterinary and Animal Sciences, Gandheli, Chhatrapati Sambhajnagar, Maharashtra

²Subject Matter Specialist, Department of Livestock Production, MGM Krishi Vigyan Kendra, Gandheli, Chhatrapati Sambhajnagar, Maharashtra

*Corresponding Author Email: sopuvet@gmail.com

Abstract

Veterinary diagnosis is moving rapidly from conventional laboratory testing toward integrated systems that combine molecular biology, digital imaging, automation, biosensors and data-driven decision support. Polymerase chain reaction (PCR), real-time PCR and emerging digital PCR methods can detect and quantify pathogen nucleic acids with high analytical sensitivity, while multiplex platforms allow several infectious agents to be examined from a single specimen. Isothermal amplification and portable point-of-care systems are bringing molecular testing closer to clinics, farms and outbreak sites where rapid decisions are essential. At the same time, automated haematology and biochemical analysers, improved immunoassays and biomarker panels are increasing the amount of clinically useful information obtained from small sample volumes. Next-generation sequencing and metagenomics can identify and characterise pathogens beyond routine targeted testing and are becoming important for outbreak investigation, microbial genomics and surveillance. Diagnostic imaging is also changing through digital radiography, ultrasonography, computed tomography and magnetic resonance imaging, with artificial intelligence increasingly used for image classification, segmentation and workflow support. Biosensors, microfluidic devices and connected monitoring systems may further shorten the interval between disease development, detection and clinical response. These technologies can improve animal welfare, treatment selection, antimicrobial stewardship and One Health surveillance, but their value depends on appropriate validation, sample quality, quality control and species-specific interpretation. This article discusses the major advanced diagnostic technologies influencing modern veterinary practice, their practical advantages and limitations, and the need to integrate new tools with clinical judgement rather than use them in isolation.

Keywords: Veterinary Diagnostics, Molecular Diagnostics, Point-Of-Care Testing, Artificial Intelligence, Biosensors, Next-Generation Sequencing, Diagnostic Imaging

How to Cite this article: Mehesare, S., & Khupse, S. M. (2026). Advanced Diagnostic Technologies in Veterinary Practice. *Future Agriculture e-magazine*, September 2026, Issue 3, pp. 204-209.

Introduction

Accurate diagnosis is the starting point of effective veterinary treatment, herd and flock health planning, disease surveillance and outbreak control. Traditional diagnosis has relied on history, clinical examination, microscopy, culture, serology, radiography and routine laboratory testing. These methods remain essential, but veterinary practice is now entering a period in which molecular assays, digital imaging, automated analysers, biosensors, sequencing and artificial

intelligence can provide information with far greater speed and depth. The change is important for both companion animals and food-producing species because diagnostic delay can lead to unnecessary treatment, continued disease transmission, production losses and avoidable antimicrobial use.

Advanced diagnostics should not be understood as a replacement for clinical judgement. A highly sensitive test can still mislead when the sample is poor, the disease stage is inappropriate or the result is interpreted without considering prevalence and clinical

signs. The real benefit of new technology comes from combining a well-collected history and examination with a test that answers a clearly defined clinical question. In this integrated approach, technology strengthens the veterinarian's decision rather than making the decision independently.

Current innovation is moving in three directions at the same time: laboratory methods are becoming more sensitive and multiplexed, sophisticated technologies are being miniaturised for point-of-care use, and digital systems are helping clinicians analyse images and large datasets. Together these developments are reducing the distance between the animal, the diagnostic laboratory and the final clinical decision.

Automation in Clinical Pathology

Automated haematology and clinical chemistry analysers have changed routine veterinary diagnostics by providing rapid, repeatable measurements from relatively small sample volumes. Modern instruments can generate complete blood counts, differential cell data, biochemical profiles, electrolyte results and other parameters within minutes. This is especially useful in emergency practice, anaesthesia, critical care and herd investigations where treatment decisions may depend on recognising anaemia, inflammation, organ dysfunction, dehydration or metabolic imbalance quickly.

Automation also creates richer datasets. Scattergrams, cell population plots and quality flags can reveal abnormalities that deserve microscopic review. However, species differences remain important. Cell size, morphology and reference intervals vary between dogs, cats, horses, cattle, small ruminants, poultry and exotic species. Therefore, analyser output should be matched to validated species settings and supported by blood-smear examination when abnormal flags or unexpected results appear. The best modern laboratory is not one that eliminates manual examination, but one that uses automation to identify where expert examination is most valuable.

PCR, qPCR and Digital Molecular Diagnostics

Polymerase chain reaction has become central to the detection of many infectious agents because it identifies pathogen nucleic acid rather than waiting for an organism to grow in culture or for an antibody response to develop. Conventional PCR is useful for targeted detection, while real-time quantitative PCR (qPCR) provides rapid amplification and fluorescent monitoring in a closed system. Multiplex qPCR can test for several pathogens or genetic targets in the same run, which is valuable when animals present with syndromes such as respiratory, enteric, reproductive or neurological disease that may have several possible causes.

Digital PCR represents a further refinement. The sample is partitioned into many small reactions and target molecules are counted statistically, allowing highly precise nucleic-acid quantification. This can be useful when pathogen load is low, when small changes in copy number matter, or when a laboratory needs a robust reference method. Nevertheless, increased analytical sensitivity does not automatically mean better clinical interpretation. Detection of nucleic acid may represent active infection, early infection, residual material after recovery or contamination. Sampling site, timing and clinical context remain essential.

Molecular diagnostics are also influencing antimicrobial stewardship. Rapid identification of a viral or specific bacterial cause can reduce empirical drug use, while resistance-gene detection can provide early information in selected settings. Culture and phenotypic antimicrobial susceptibility testing remain important because the presence of a resistance gene does not always predict the complete behaviour of an organism. A combined molecular and culture-based approach often provides the most clinically useful picture.

Isothermal Amplification and Point-of-Care Testing


One limitation of conventional PCR is its dependence on thermal cycling equipment and trained laboratory personnel. Isothermal amplification technologies, including loop-mediated isothermal amplification (LAMP) and

recombinase-based methods, amplify nucleic acid at a constant temperature. This simplifies equipment needs and creates opportunities for testing at clinics, farms, field camps and outbreak sites. Many assays can produce results in less than an hour and can be linked with visual, fluorescent or lateral-flow readouts.

Point-of-care tests are particularly valuable when immediate action is required. A veterinarian facing a contagious disease can isolate animals, advise movement restrictions, begin supportive treatment or collect confirmatory samples without waiting several days for a distant laboratory. Rapid tests are also useful in resource-limited areas. Their main challenge is quality assurance: a fast result is only helpful if the assay has appropriate sensitivity, specificity and validation for the target species, specimen and intended use. Positive and negative controls, correct storage and clear interpretation rules remain essential even for simple field tests.

Immunodiagnosics and Biomarker-Based Testing


Serological and antigen-detection methods continue to be major tools in veterinary diagnosis. Enzyme-linked immunosorbent assays, lateral-flow immunoassays and other antibody or antigen platforms can support surveillance, vaccination monitoring and diagnosis of infectious diseases. Newer formats use recombinant antigens, monoclonal antibodies and improved signal systems to increase specificity and reduce background reactions. Multiplex immunoassays can measure several antibodies or biomarkers from one sample, conserving time and sample volume.



Biomarker research is expanding diagnosis beyond the simple presence or absence of a pathogen. Acute-phase proteins, cardiac biomarkers, endocrine markers, inflammatory mediators and tumour-associated molecules can help assess disease severity, organ injury and treatment response. The clinical value of a biomarker depends on validated reference ranges and on understanding what biological process it represents. Biomarker panels may be more informative than a single measurement because diseases often affect several pathways simultaneously.

Next-Generation Sequencing and Metagenomics

Next-generation sequencing (NGS) can analyse millions of DNA or RNA fragments in parallel. In veterinary infectious disease work it can identify and characterise pathogens, compare outbreak strains and detect genetic changes that may influence epidemiology, virulence or resistance. Whole-genome sequencing is especially powerful after an organism has been isolated, while metagenomic sequencing can examine all genetic material in a sample and may detect unexpected or difficult-to-culture agents.



Metagenomics is attractive when routine testing is negative but an infectious cause is still suspected. It can also be used to study microbiomes and understand how microbial communities change with disease, diet, antimicrobial exposure or management. The challenge is interpretation. Sequencing can detect harmless commensals, environmental contaminants and nucleic acid from organisms that are not responsible for the clinical problem. Bioinformatics, contamination control and

carefully selected clinical metadata are therefore as important as the sequencing instrument itself.

Advanced Imaging and Artificial Intelligence

Veterinary diagnostic imaging has progressed from conventional radiography to high-resolution ultrasound, computed tomography (CT), magnetic resonance imaging (MRI), endoscopy and other specialised modalities. These technologies allow clinicians to evaluate anatomy and function non-invasively and have transformed orthopaedics, neurology, cardiology, oncology and emergency care. Portable digital radiography and ultrasound have also brought advanced imaging to equine, farm and field practice.

Artificial intelligence (AI) is increasingly being studied as a decision-support tool for image analysis. Machine-learning models can be trained to classify radiographs, segment structures, measure lesions and identify patterns that may be difficult to quantify manually. Recent veterinary literature shows growing applications across radiography, CT, MRI and ultrasound. AI may assist with workflow, triage and quality control, particularly where access to specialist radiologists is limited.

However, an AI result is not automatically a diagnosis. Models can perform poorly when an image comes from a different machine, species, breed, position or clinical population than the data used for training. Incorrect or inconsistent labels in training datasets can also become embedded in the model. Veterinary AI therefore requires external validation, transparent reporting and clinician oversight. The safest role is to provide a second layer of analysis while the veterinarian remains responsible for integrating the image with the animal's history and examination.

Biosensors, Microfluidics and Connected Diagnostics

Biosensors convert a biological interaction into a measurable signal. Electrochemical, optical and other sensor platforms are being developed for infectious agents, metabolites, hormones and markers of inflammation. When combined with microfluidics, very small volumes of blood, saliva, milk or other fluids

can be processed through compact devices. The long-term goal is a lab-on-a-chip system capable of sample preparation, reaction, detection and digital reporting in one portable unit.

Connected diagnostic devices can also link testing with electronic medical records and surveillance systems. Wearable sensors used for temperature, activity, rumination, heart rate or behaviour may identify animals that require closer examination before obvious clinical signs develop. In dairy and livestock systems, this creates the possibility of moving from periodic health checks to continuous risk monitoring. Such systems are most valuable when alerts are clinically meaningful and false alarms are controlled; otherwise large amounts of data can increase workload rather than improve care.

Table 1. Major advanced diagnostic technologies used in veterinary practice.

Technology	Main veterinary use
qPCR / digital PCR	Sensitive detection or quantification of pathogen nucleic acid in validated specimens.
Isothermal amplification	Rapid field or near-patient detection of selected DNA/RNA targets with minimal equipment.
ELISA / rapid immunoassay	Detection of antibodies, antigens or selected biomarkers for screening and clinical support.
NGS / metagenomics	Broad investigation of unexpected, mixed or genetically variable infectious agents.
Advanced imaging	Ultrasound, CT or MRI localize structural or functional abnormalities and guide clinical decisions.
Biosensors / microfluidics	Portable, low-volume detection of biological signals with potential point-of-care connectivity.

Diagnostic Integration, One Health and Surveillance

Advanced diagnostics have an important population-level role. Rapid and accurately geolocated results can strengthen surveillance for transboundary and zoonotic diseases, support tracing during outbreaks and provide earlier warning of unusual disease patterns. Genomic information can help determine whether cases are linked, while digital dashboards can combine laboratory, farm and movement data. This is directly relevant to One Health because many pathogens circulate between animals, people and the environment.

The same principle applies inside a veterinary hospital or farm. Diagnostic information becomes more useful when results from pathology, imaging, microbiology and

clinical records can be viewed together. Integrated interpretation may reveal relationships that are less obvious when each test is considered separately. Interoperable data systems and standard terminology will therefore become increasingly important as the number and complexity of tests grow.

Quality Assurance and Practical Limitations

Table 2. Sample-to-decision steps for reliable veterinary diagnosis.

Stage	Good practice
Clinical question	Define the disease problem, animal population and decision the test is expected to support.
Sample collection	Collect the correct specimen at the appropriate disease stage with suitable technique and identification.
Transport and storage	Maintain required time, temperature and chain of identity from animal to laboratory.
Test selection	Match assay validation, sensitivity and specificity to the species, pathogen and intended diagnostic purpose.
Interpretation	Combine the laboratory result with clinical findings, exposure history, prevalence and test limitations.

Every diagnostic technology has limitations. Pre-analytical errors such as incorrect sample type, delayed transport, contamination, haemolysis or poor storage can invalidate even the most sophisticated assay. Analytical quality depends on calibration, controls, reagent performance and validated procedures. Post-analytical quality depends on correct interpretation and communication. A technically correct laboratory result can still cause harm if it is reported without appropriate clinical context.

Cost and accessibility also influence adoption. High-end sequencing, CT and MRI systems require infrastructure and trained specialists, while some rapid platforms have higher per-test costs than central laboratory methods. Veterinary practices therefore need to choose technology according to case volume, disease risk and the consequences of delayed diagnosis. Referral laboratories will remain essential even as more testing moves closer to the patient.

Validation is especially important when a human diagnostic platform is adapted for veterinary use. Species-specific differences may affect antibodies, biomarkers, reference intervals and sample matrices. International standards emphasise that diagnostic assays should be validated for their intended purpose.

Sensitivity and specificity are not fixed properties in every situation; performance may change with population, disease stage and specimen type. Good diagnostics depend on knowing both what a test can detect and what it cannot.

Conclusion

Advanced diagnostic technologies are reshaping veterinary practice by bringing laboratory precision closer to the animal and by allowing clinicians to obtain more information from smaller samples and shorter time intervals. PCR and digital molecular assays, isothermal amplification, improved immunodiagnosics, sequencing, advanced imaging, artificial intelligence, biosensors and connected monitoring systems each address different parts of the diagnostic process.

The central principle remains unchanged: technology must serve sound clinical reasoning. Reliable sampling, validated methods, quality control, species-specific interpretation and communication with animal owners or producers remain essential. When these foundations are maintained, new diagnostic tools can support earlier disease recognition, more targeted treatment, responsible antimicrobial use, stronger surveillance and better animal health outcomes.

References

- Barillaro, G., Minniti, S., Mezzatesta, S., Macri, F., Torino, C., & Vilasi, A. D. (2026). Artificial intelligence in veterinary diagnostic imaging: A narrative review of the state of the art and its applications. *The Veterinary Journal*, 318, 106762. <https://doi.org/10.1016/j.tvjl.2026.106762>
- Bora, M., Manu, M., Mathew, D. D., Das, H., Bora, D. P., & Barman, N. N. (2022). Point of care diagnostics and non-invasive sampling strategy: A review on major advances in veterinary diagnostics. *Acta Veterinaria Brno*, 91, 17-34. <https://doi.org/10.2754/avb202291010017>
- Burti, S., Banzato, T., Coghlan, S., Wodzinski, M., Bendazzoli, M., & Zotti, A. (2024). Artificial intelligence in veterinary diagnostic imaging: Perspectives and limitations. *Research in Veterinary Science*, 175, 105317. <https://doi.org/10.1016/j.rvsc.2024.105317>
- Hennessey, E., DiFazio, M., Hennessey, R., & Cassel, N. (2022). Artificial intelligence in veterinary diagnostic imaging: A literature review. *Veterinary Radiology & Ultrasound*, 63(Suppl. 1), 851-870. <https://doi.org/10.1111/vru.13163>
- Nealon, N. J. (2026). Next-generation sequencing in companion animal practice for infectious disease



- diagnostics and characterizing normal microbiomes. *Veterinary Clinics of North America: Small Animal Practice*. <https://doi.org/10.1016/j.cvsm.2026.06.005>
- Sahoo, P. R., Sethy, K., Mohapatra, S., & Panda, D. (2016). Loop mediated isothermal amplification: An innovative gene amplification technique for animal diseases. *Veterinary World*, 9(5), 465-469.
- Staji, H., & Angoshtan, S. F. (2026). Biosensors for the detection of arthropod-borne veterinary viruses: A comprehensive review. *Veterinary and Animal Science*, 32, 100577. <https://doi.org/10.1016/j.vas.2026.100577>
- Tooryan, F., Enyertornye, B., Choudhury, F. J., & Velayudhan, B. T. (2026). Recent advances in isothermal amplification techniques for detection of animal diseases. *Frontiers in Veterinary Science*, 13, 1819190. <https://doi.org/10.3389/fvets.2026.1819190>
- World Organisation for Animal Health (WOAH). (2024). *Manual of Diagnostic Tests and Vaccines for Terrestrial Animals* (13th ed.). Paris: WOAH.