

Conservation Agriculture and Carbon Sequestration Strategies

Banhisikha Roy¹, Saikat Das², Ardhendu Swain³ and Deep Prabha^{*4}

^{1,2,3,4}Department of Soil Science and Agricultural Chemistry, Palli Siksha Bhavana (Institute of Agriculture), Visva-Bharati, Sriniketan-731236, West Bengal

***Corresponding Author Email:** deeprabha.rs.solscie@visva-bharati.ac.in

ORCID ID: 0000-0002-0271-6622

Abstract

Conservation agriculture is based on minimum soil disturbance, permanent soil cover and diversified crop rotations. These principles can improve soil structure, reduce erosion, conserve water and influence the storage of organic carbon in agricultural soils. Carbon sequestration occurs when carbon inputs from crop residues, roots, cover crops or organic amendments exceed carbon losses through decomposition and erosion. Reduced tillage can slow physical disturbance and help protect soil aggregates, while residue retention and diverse rotations increase organic inputs. The response is not uniform in every soil. Climate, texture, initial soil carbon, crop productivity, residue use and sampling depth strongly influence measured changes. In some systems, reduced tillage redistributes carbon toward the soil surface without producing the same increase at deeper depths, so assessment should consider the full profile relevant to management. Conservation agriculture also interacts with nitrogen cycling and may affect nitrous oxide emissions, which means climate benefits should be evaluated using a broader greenhouse-gas perspective. Carbon sequestration should not be treated as a substitute for reducing fossil-fuel emissions, but it can contribute to soil restoration and climate mitigation while supporting productivity. Long-term adoption, accurate measurement and locally adapted rotations are essential for credible outcomes.

Keywords: Conservation Agriculture, Soil Organic Carbon, Carbon Sequestration, Residue Retention, Reduced Tillage, Crop Rotation

How to Cite this article: Roy, B., Das, S., Prabha, D., & Swain, A. (2026). Conservation Agriculture and Carbon Sequestration Strategies. *Future Agriculture e-magazine*, September 2026, Issue 3, pp. 288-294.

Introduction

Agricultural soils contain large stores of carbon, but cultivation can reduce soil organic matter through erosion, residue removal and accelerated decomposition. Conservation agriculture aims to maintain a more protected soil environment while sustaining crop production. Practical implementation of introduction should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to improving soil protection, biological inputs and crop diversity while evaluating carbon storage as a long-term co-benefit. A practical plan is to establish the current tillage, residue, rotation and soil-carbon baseline before changing management so later comparisons are

meaningful. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include soil cover, erosion, residue return, crop yield, soil organic carbon, fuel use and input efficiency. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that carbon gains are gradual and reversible, so short trials or surface-only sampling can give misleading conclusions. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be


maintained and to produce reliable long-term benefit.

Its three commonly recognised principles are minimum mechanical disturbance, continuous soil cover and crop diversification. Carbon sequestration is a possible co-benefit when these practices increase organic inputs and reduce losses over a sufficiently long period. The value of introduction also depends on integration with the wider production system. Management should connect improving soil protection, biological inputs and crop diversity while evaluating carbon storage as a long-term co-benefit with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for soil cover, erosion, residue return, crop yield, soil organic carbon, fuel use and input efficiency. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because carbon gains are gradual and reversible, so short trials or surface-only sampling can give misleading conclusions, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Soil Organic Carbon Dynamics

Soil carbon is supplied mainly by roots, crop residues, manures and other organic materials. Microorganisms transform these inputs, and some carbon is stabilised through association with minerals or protection within aggregates. Temperature, moisture, aeration and soil texture influence decomposition. Practical implementation of soil organic carbon dynamics should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to balancing carbon inputs from roots and residues against decomposition, erosion and other losses. A practical plan is to consider crop productivity, root distribution, texture, moisture and mineral interactions when interpreting

changes in soil carbon. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include organic carbon by depth, bulk density, residue inputs, root biomass, aggregate stability and seasonal moisture. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that soils approach a new equilibrium and do not accumulate carbon indefinitely under any single practice. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.



A management change does not create unlimited carbon storage. Soils approach a new equilibrium over time, and gains can be reversed if practices are discontinued. Baseline carbon and crop productivity therefore determine the potential for additional sequestration. The value of soil organic carbon dynamics also depends on integration with the wider production system. Management should connect balancing carbon inputs from roots and residues against decomposition, erosion and other losses with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for organic carbon by depth, bulk density, residue inputs, root biomass, aggregate stability and

seasonal moisture. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because soils approach a new equilibrium and do not accumulate carbon indefinitely under any single practice, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Minimum Tillage and Soil Disturbance

Reduced or zero tillage limits mechanical disruption of soil aggregates and can reduce fuel use. Surface residues also protect soil from raindrop impact and may increase infiltration. These effects are particularly valuable where erosion is a major source of soil-carbon loss. Practical implementation of minimum tillage and soil disturbance should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to reducing aggregate disruption, erosion and fuel use while maintaining reliable crop establishment.



A practical plan is to adapt seeding equipment, traffic patterns and weed management to local soil texture, drainage and residue load rather than assuming one tillage intensity fits all fields. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include surface cover, fuel use, penetration resistance, bulk density, emergence, weed pressure and carbon by depth. Trends are more useful than isolated readings because

season, production stage and local conditions alter expected performance. One limitation is that reduced tillage can redistribute carbon toward the surface and may create compaction or weed challenges if other management is not adjusted. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Carbon responses depend on sampling depth. No-till systems often show higher carbon near the surface, while whole-profile changes may be smaller. Compaction, weed management and crop establishment must also be managed carefully during transition. The value of minimum tillage and soil disturbance also depends on integration with the wider production system. Management should connect reducing aggregate disruption, erosion and fuel use while maintaining reliable crop establishment with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for surface cover, fuel use, penetration resistance, bulk density, emergence, weed pressure and carbon by depth. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because reduced tillage can redistribute carbon toward the surface and may create compaction or weed challenges if other management is not adjusted, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Table 1. Conservation agriculture practices influencing soil carbon.

Practice	Carbon-related mechanism
Reduced / zero tillage	Limits disturbance and can reduce erosion and aggregate disruption.
Residue retention	Adds organic carbon and protects the soil surface.
Cover crops	Increase living-root duration and off-season biomass inputs.

Practice	Carbon-related mechanism
Diverse rotations	Provide varied root systems and residue qualities and may improve productivity.
Organic amendments	Add external organic matter and nutrients when used responsibly.
Erosion control	Retains carbon-rich topsoil that would otherwise be lost from the field.

Residue Retention and Soil Cover

Crop residues provide carbon and nutrients and reduce evaporation and erosion. Maintaining cover moderates soil temperature and provides substrate for soil organisms. In dry regions, improved water conservation can also increase crop biomass and root inputs. Practical implementation of residue retention and soil cover should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to protecting the surface and returning carbon and nutrients through crop residues, mulch and cover crops. A practical plan is to quantify competing residue uses, retain as much cover as practical and manage residues so they do not interfere with sowing or create disease and pest problems. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include percentage ground cover, residue biomass, erosion, soil moisture, temperature, infiltration and subsequent crop establishment. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that complete retention may be unrealistic in mixed crop-livestock systems, so partial retention and alternative fodder sources may be more sustainable. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Residue has competing uses for fodder, fuel or bedding, so complete retention may not be practical. Partial retention, cover crops or organic amendments can be used to balance soil protection with farm needs. The value of residue retention and soil cover also depends on integration with the wider production system.

Management should connect protecting the surface and returning carbon and nutrients through crop residues, mulch and cover crops with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for percentage ground cover, residue biomass, erosion, soil moisture, temperature, infiltration and subsequent crop establishment. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because complete retention may be unrealistic in mixed crop-livestock systems, so partial retention and alternative fodder sources may be more sustainable, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Crop Rotation and Cover Crops

Diverse rotations produce different rooting patterns and residue qualities. Legumes can contribute biologically fixed nitrogen, while deep-rooted crops may place carbon at greater depth. Cover crops add biomass during periods when soil would otherwise remain bare. Practical implementation of crop rotation and cover crops should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to increasing rooting diversity, biological nitrogen inputs and year-round carbon flow to soil. A practical plan is to choose rotations and cover species that fit rainfall, planting windows, market needs and water availability, and terminate covers at an appropriate stage. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include rotation diversity, biomass, root growth, nitrogen contribution, soil water, yield stability and pest pressure. Trends are more useful than isolated readings because season, production

stage and local conditions alter expected performance. One limitation is that cover crops require seed, labour and water, and poorly timed termination can compete with the following cash crop. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Rotation also reduces pest and disease pressure and can spread labour and weather risk. The carbon benefit depends on maintaining total biomass production; a diverse rotation that sharply reduces annual carbon input may not increase soil carbon. The value of crop rotation and cover crops also depends on integration with the wider production system. Management should connect increasing rooting diversity, biological nitrogen inputs and year-round carbon flow to soil with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for rotation diversity, biomass, root growth, nitrogen contribution, soil water, yield stability and pest pressure. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because cover crops require seed, labour and water, and poorly timed termination can compete with the following cash crop, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Nutrient and Water Management

Carbon sequestration requires plant growth, so severe nutrient or water limitation restricts the amount of biomass returned to soil. Balanced fertilisation and efficient irrigation can increase residue and root inputs, although excessive nitrogen may increase nitrous oxide emissions. Practical implementation of nutrient and water management should start with a clear

baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to supporting biomass production and carbon inputs without creating nutrient loss or greenhouse-gas trade-offs. A practical plan is to combine soil testing, balanced fertiliser, organic amendments, residue management and water conservation to remove avoidable limits on crop growth. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include nutrient balance, water-use efficiency, biomass return, yield, nitrous oxide risk, leaching and soil organic carbon. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that carbon sequestration claims are weak when gains depend on excessive fertiliser, irrigation or off-farm organic inputs that shift impacts elsewhere. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Integrated nutrient management combines fertiliser with organic sources where available. Water conservation through mulch and reduced disturbance can improve drought resilience and support more stable carbon inputs across seasons. The value of nutrient and water management also depends on integration with the wider production system. Management should connect supporting biomass production and carbon inputs without creating nutrient loss or greenhouse-gas trade-offs with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for nutrient balance, water-use efficiency, biomass return, yield, nitrous oxide risk, leaching and soil organic carbon. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because carbon sequestration claims are weak when gains

depend on excessive fertiliser, irrigation or off-farm organic inputs that shift impacts elsewhere, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Table 2. Key considerations for measuring soil carbon change.

Consideration	Good practice
Baseline	Measure before or at the start of management change where possible.
Sampling depth	Use consistent depths and include deeper layers when relevant.
Bulk density	Measure so carbon concentration can be converted to stock per area.
Spatial variability	Use replicated or georeferenced sampling locations.
Time interval	Allow sufficient time because soil carbon changes slowly.
Greenhouse-gas balance	Consider nitrous oxide and energy use when making climate claims.

Measurement and Verification

Soil organic carbon should be measured using consistent sampling depth, bulk density and laboratory methods. Because carbon changes slowly and soil is spatially variable, short-term measurements can be misleading. Repeated sampling at fixed locations or robust field designs improve confidence. Practical implementation of measurement and verification should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil carbon sequestration, priority should be given to measuring small changes in a large and variable soil carbon pool with consistent methods and adequate time. A practical plan is to use fixed sampling locations or sound designs, consistent depths, bulk-density measurements and accredited laboratory methods, and document management history. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include carbon concentration, carbon stock, equivalent soil mass where appropriate, spatial variability, sampling error and change through time. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that changes in bulk density or sampling depth

can create apparent gains or losses that are not real changes in total soil carbon. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Equivalent soil mass approaches may be useful where bulk density changes after tillage conversion. Carbon accounting should also consider changes in fuel use, fertiliser and nitrous oxide where climate mitigation is the goal. The value of measurement and verification also depends on integration with the wider production system. Management should connect measuring small changes in a large and variable soil carbon pool with consistent methods and adequate time with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for carbon concentration, carbon stock, equivalent soil mass where appropriate, spatial variability, sampling error and change through time. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because changes in bulk density or sampling depth can create apparent gains or losses that are not real changes in total soil carbon, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Co-Benefits and Trade-Offs

Conservation agriculture can reduce erosion, improve infiltration, increase aggregate stability and support biological activity. These improvements may increase yield stability even when soil-carbon gains are modest. Practical implementation of co-benefits and trade-offs should start with a clear baseline and a defined management objective. For conservation agriculture and credible soil

carbon sequestration, priority should be given to evaluating erosion control, infiltration, biodiversity, fuel use and yield stability alongside carbon storage. A practical plan is to compare multiple outcomes and transition costs so conservation practices are selected for whole-system value rather than carbon alone. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include yield, profitability, erosion, infiltration, labour, herbicide use, biodiversity indicators, fuel and carbon stock. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that some systems experience temporary yield or weed-management challenges, and benefits vary with climate, soil and management quality. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Trade-offs include residue competition, specialised seeding equipment, weed shifts and variable yield response during transition. Local experimentation and farmer support are therefore important for long-term adoption. The value of co-benefits and trade-offs also depends on integration with the wider production system. Management should connect evaluating erosion control, infiltration, biodiversity, fuel use and yield stability alongside carbon storage with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for yield, profitability, erosion, infiltration, labour, herbicide use, biodiversity indicators, fuel and carbon stock. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because some systems experience temporary yield or weed-management challenges, and benefits vary with climate, soil and management quality, weak results should first

prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Conclusion

Conservation agriculture can support soil carbon sequestration when reduced disturbance is combined with continuous organic inputs and diverse cropping. Residue retention, cover crops, productive rotations and balanced nutrient management are central to building soil organic matter.

Carbon gains must be measured carefully across appropriate soil depths and time periods. The strongest case for conservation agriculture is its combination of soil protection, resilience and potential climate benefits rather than carbon storage alone.

References

- Kassam, A., Friedrich, T., Shaxson, F., & Pretty, J. (2009). The spread of conservation agriculture: Justification, sustainability and uptake. *International Journal of Agricultural Sustainability*, 7, 292-320.
- Lal, R. (2004). Soil carbon sequestration impacts on global climate change and food security. *Science*, 304, 1623-1627.
- Pittelkow, C. M., Liang, X., Linquist, B. A., et al. (2015). Productivity limits and potentials of the principles of conservation agriculture. *Nature*, 517, 365-368.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., & Smith, P. (2016). Climate-smart soils. *Nature*, 532, 49-57.
- Lehmann, J., Hansel, C. M., Kaiser, C., et al. (2020). Persistence of soil organic carbon caused by functional complexity. *Nature Geoscience*, 13, 529-534.
- FAO. (2017). *Conservation Agriculture*. Food and Agriculture Organization of the United Nations, Rome.