

Soil Microbiome and Its Role in Sustainable Crop Production

Manjot Singh^{*1} and Shagun²

¹Ph.D., Soil Science; Lecturer, Surender Kaur Memorial Agriculture College, 24 BB, Padampur, Sri Ganganagar, Rajasthan, India

²M.Sc., Department of Vegetable Science, Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India

*Corresponding Author Email: manjotchinzer97@gmail.com

Abstract

The soil microbiome includes bacteria, fungi, archaea, protists and other microorganisms that interact with plant roots and organic matter. These organisms regulate decomposition, nutrient cycling, soil aggregation and many biological processes that influence crop growth. The rhizosphere is particularly active because roots release carbon compounds that attract and select microbial communities. Some microbes fix nitrogen, solubilise phosphorus, produce growth-promoting compounds or form mycorrhizal associations that improve nutrient and water acquisition. Others can suppress plant pathogens through competition, antibiosis or stimulation of plant defence. The microbiome is also influenced by tillage, crop rotation, fertiliser, pesticides, irrigation, organic amendments and soil pH. Management that maintains organic inputs and diverse plant cover generally supports a more active biological community, although the response depends on soil and climate. Commercial biofertilisers and microbial inoculants can be useful when effective strains are matched to the crop and environment, but field performance is often less predictable than laboratory results because introduced organisms must compete with native communities. New molecular tools are improving understanding of microbial diversity and function. Sustainable crop production can benefit from microbiome-aware management, but it should integrate biology with sound soil fertility, water and disease management rather than treating microbes as a stand-alone solution.

Keywords: Soil Microbiome, Rhizosphere, Nutrient Cycling, Mycorrhizae, Biofertiliser, Sustainable Agriculture

How to Cite this article: Singh, M. (2026). Soil Microbiome and Its Role in Sustainable Crop Production. *Future Agriculture e-magazine*, September 2026, Issue 3, pp. 295-302.

Introduction

A gram of healthy soil can contain enormous numbers of microorganisms representing many functional groups. Although most are not visible, their combined activity influences how residues decompose, nutrients become available and soil structure develops. Practical implementation of introduction should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to recognising soil microorganisms as an interacting community that influences decomposition, nutrient availability, structure and root health. A practical plan is to manage the rhizosphere through crop diversity, organic inputs, balanced fertility and suitable moisture rather than trying to control individual microbes in isolation. Changes should be introduced consistently so


their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include organic matter, root growth, nutrient availability, respiration or other biological indicators, disease and crop performance. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that microbial abundance alone does not prove beneficial function because community activity depends on soil and plant context. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Plant roots create a specialised zone called the rhizosphere. Root exudates provide energy-rich compounds that shape microbial communities, while microbes can influence plant nutrition, growth and defence. This two-way interaction is central to the soil microbiome. The value of introduction also depends on integration with the wider production system. Management should connect recognising soil microorganisms as an interacting community that influences decomposition, nutrient availability, structure and root health with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for organic matter, root growth, nutrient availability, respiration or other biological indicators, disease and crop performance. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because microbial abundance alone does not prove beneficial function because community activity depends on soil and plant context, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Microbial Diversity and Soil Function

Bacteria and fungi dominate many decomposition and nutrient-cycling processes, while archaea participate in nitrogen transformations and protists influence microbial food webs. The importance of a Practical implementation of microbial diversity and soil function should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to supporting bacterial, fungal, archaeal and protist functions that contribute to nutrient cycling and soil structure. A practical plan is to maintain diverse carbon substrates, avoid repeated severe disturbance and consider pH, moisture and aeration as major filters of

microbial activity. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include microbial biomass or activity, aggregation, decomposition, soil pH, moisture, residue turnover and crop response. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that greater diversity is not automatically better if the functional groups needed for a particular process are absent or inactive. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.



group depends on substrate, moisture, pH and oxygen conditions. The value of microbial diversity and soil function also depends on integration with the wider production system. Management should connect supporting bacterial, fungal, archaeal and protist functions that contribute to nutrient cycling and soil structure with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for microbial biomass or activity, aggregation, decomposition, soil pH, moisture, residue turnover and crop response. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because greater diversity is not

automatically better if the functional groups needed for a particular process are absent or inactive, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Diversity alone does not guarantee a beneficial function. Sustainable management aims to create conditions that support organic-matter turnover, aggregation and nutrient retention while avoiding disturbance that repeatedly reduces biological activity.

Nitrogen, Phosphorus and Nutrient Cycling

Symbiotic bacteria in legume nodules convert atmospheric nitrogen into forms that plants can use. Free-living and associative nitrogen fixers may also contribute smaller amounts. Other microorganisms mineralise organic nitrogen and influence nitrification and denitrification. Practical implementation of nitrogen, phosphorus and nutrient cycling should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to using biological nitrogen fixation, mineralisation, nitrification, phosphorus mobilisation and mycorrhizal acquisition within balanced fertility programmes. A practical plan is to match legume inoculation and nutrient management to crop need, soil test results and local environmental conditions. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include nodulation, plant nitrogen status, available phosphorus, fertiliser response, nutrient-use efficiency and yield. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that microbial processes can also contribute to nitrogen loss through denitrification or immobilisation, so biological activity must be interpreted in context. Therefore records should be reviewed regularly and important deviations checked by the

relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.


Phosphate-solubilising bacteria and fungi can increase availability of some poorly soluble phosphorus compounds. Mycorrhizal fungi extend the effective absorbing network of roots and are especially important for phosphorus acquisition under low-input conditions. The value of nitrogen, phosphorus and nutrient cycling also depends on integration with the wider production system. Management should connect using biological nitrogen fixation, mineralisation, nitrification, phosphorus mobilisation and mycorrhizal acquisition within balanced fertility programmes with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for nodulation, plant nitrogen status, available phosphorus, fertiliser response, nutrient-use efficiency and yield. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because microbial processes can also contribute to nitrogen loss through denitrification or immobilisation, so biological activity must be interpreted in context, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Table 1. Important soil microbial groups and major functions.

Microbial group	Examples of functions
Rhizobia	Symbiotic nitrogen fixation in legumes.
Plant-growth-promoting rhizobacteria	Nutrient mobilisation, hormone effects and induced plant defence.
Arbuscular mycorrhizal fungi	Improved phosphorus and water acquisition and contribution to aggregation.
Decomposer fungi and bacteria	Breakdown of residues and release or immobilisation of nutrients.
Nitrifiers / denitrifiers	Transform soil nitrogen and influence nitrogen-use efficiency and gaseous losses.
Antagonistic microbes	Compete with or inhibit selected soil-borne plant pathogens.

Mycorrhizae and Root Symbiosis

Arbuscular mycorrhizal fungi colonise roots of many crops and form hyphal networks in soil. These networks can improve access to phosphorus and water and may influence soil aggregation. Benefits are usually greater where nutrient availability is moderate rather than excessive. Practical implementation of mycorrhizae and root symbiosis should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to maintaining beneficial root-fungal associations that extend nutrient and water acquisition beyond the immediate root surface. A practical plan is to reduce unnecessary disruption, use compatible crop rotations and avoid nutrient regimes that eliminate the plant's incentive to maintain symbiosis.



Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include root colonisation where assessed, phosphorus uptake, drought response, root growth, crop vigour and yield. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that mycorrhizal benefits vary among crops, soils and fertility levels and are not guaranteed by the presence of fungal inoculum alone. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be

maintained and to produce reliable long-term benefit.

Intensive tillage can disrupt fungal networks, while long fallows and non-host crops can reduce inoculum. Crop rotation and reduced disturbance may therefore influence the strength of mycorrhizal associations. The value of mycorrhizae and root symbiosis also depends on integration with the wider production system. Management should connect maintaining beneficial root-fungal associations that extend nutrient and water acquisition beyond the immediate root surface with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for root colonisation where assessed, phosphorus uptake, drought response, root growth, crop vigour and yield. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because mycorrhizal benefits vary among crops, soils and fertility levels and are not guaranteed by the presence of fungal inoculum alone, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Disease Suppression and Plant Defence

Some soils naturally suppress particular pathogens because resident microbes compete for nutrients, occupy infection sites or produce inhibitory compounds. Beneficial rhizobacteria can also stimulate plant defence responses. Practical implementation of disease suppression and plant defence should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to encouraging microbial competition, antibiosis and induced plant responses that reduce opportunities for soil-borne pathogens. A practical plan is to combine diverse rotations, healthy organic inputs, sanitation and resistant varieties so biological suppression

complements established disease management. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include disease incidence, pathogen populations, root health, crop stand, amendment response and yield. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that suppressive effects are often community-level and site-specific, making single-organism claims difficult to reproduce across fields. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Disease suppression is usually a community property rather than the result of one organism. Organic amendments and rotations may shift microbial communities, but poorly decomposed material can also introduce pathogens or create temporary nutrient imbalance. The value of disease suppression and plant defence also depends on integration with the wider production system. Management should connect encouraging microbial competition, antibiosis and induced plant responses that reduce opportunities for soil-borne pathogens with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for disease incidence, pathogen populations, root health, crop stand, amendment response and yield. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because suppressive effects are often community-level and site-specific, making single-organism claims difficult to reproduce across fields, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process

allows timing, thresholds and resource use to be refined as experience accumulates.

Microbiome and Abiotic Stress Tolerance

Root-associated microbes can influence plant response to drought, salinity and nutrient stress by changing root growth, hormone signalling or nutrient acquisition. Research on these interactions is expanding because climate stress is becoming more important in crop production. Practical implementation of microbiome and abiotic stress tolerance should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to understanding how root-associated microbes may influence drought, salinity and nutrient-stress responses through root growth and signalling. A practical plan is to test promising practices under local field stress, maintain realistic controls and separate microbial effects from changes in water or fertility. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include plant water status, salinity response, root traits, survival, biomass, nutrient uptake and yield stability. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that results from controlled environments may not persist in field soils where introduced organisms compete with established communities. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Field effects are variable. A strain that improves stress tolerance in controlled conditions may not establish in a different soil. Local testing is therefore necessary before broad recommendation. The value of microbiome and abiotic stress tolerance also depends on integration with the wider production system. Management should connect understanding how root-associated microbes may influence drought, salinity and

nutrient-stress responses through root growth and signalling with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for plant water status, salinity response, root traits, survival, biomass, nutrient uptake and yield stability. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because results from controlled environments may not persist in field soils where introduced organisms compete with established communities, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Table 2. Farm practices that can influence the soil microbiome.

Practice	Likely microbiome effect
Diverse crop rotation	Broadens root exudates and residue inputs across seasons.
Cover cropping	Maintains living roots and microbial activity between cash crops.
Organic amendments	Adds carbon substrates and nutrients for microbial growth.
Reduced tillage	Reduces physical disruption of aggregates and fungal networks.
Balanced fertilisation	Supports plant growth without extreme nutrient or pH effects.
Good drainage / irrigation	Maintains moisture and oxygen conditions suitable for desired processes.

Management Practices that Shape the Microbiome

Crop diversity, cover crops and organic inputs provide a wider range of substrates for soil organisms. Reduced tillage may protect fungal networks and aggregates. Balanced fertilisation supports plant growth and root carbon inputs, while extreme nutrient application or soil acidity can shift communities. Practical implementation of management practices that shape the microbiome should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to using


rotations, cover crops, organic amendments, tillage, irrigation and pesticides to create a biologically favourable soil environment. A practical plan is to evaluate management as a package over several seasons because microbial communities respond to cumulative changes in substrate and disturbance. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include soil cover, residue inputs, moisture, pH, microbial activity, disease, nutrient cycling and crop performance. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that heavy disturbance, repeated bare fallow or inappropriate chemical use can reduce biological function even when microbial products are added. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Pesticides should be used according to need and label because non-target effects vary among compounds and microbial groups. Irrigation and drainage also shape the microbiome by controlling oxygen and moisture. The value of management practices that shape the microbiome also depends on integration with the wider production system. Management should connect using rotations, cover crops, organic amendments, tillage, irrigation and pesticides to create a biologically favourable soil environment with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for soil cover, residue inputs, moisture, pH, microbial activity, disease, nutrient cycling and crop performance. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because heavy disturbance, repeated bare fallow or inappropriate chemical use can reduce biological function even when microbial

products are added, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Biofertilisers and Microbial Inoculants

Rhizobium, Azotobacter, Azospirillum, phosphate-solubilising microorganisms and mycorrhizal inoculants are commonly promoted as biofertilisers. Their value depends on strain quality, viable cell count, storage, application method and compatibility with the crop. Practical implementation of biofertilisers and microbial inoculants should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to using verified strains of Rhizobium, Azotobacter, Azospirillum, phosphate-solubilising organisms or mycorrhizae where crop and field conditions are suitable.



A practical plan is to select quality products, protect viability during storage, apply at the recommended time and compare performance with standard nutrient management. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include viable count where available, nodulation, establishment, nutrient uptake, yield response, cost and consistency across fields. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that inoculants should not

replace soil testing or adequate fertility, and poor-quality or poorly matched products may show little field benefit. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Inoculants should complement rather than replace basic soil testing and nutrient management. Farmers should use products with quality assurance and realistic claims, especially because performance can vary with temperature, soil pH and native microbial competition. The value of biofertilisers and microbial inoculants also depends on integration with the wider production system. Management should connect using verified strains of Rhizobium, Azotobacter, Azospirillum, phosphate-solubilising organisms or mycorrhizae where crop and field conditions are suitable with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for viable count where available, nodulation, establishment, nutrient uptake, yield response, cost and consistency across fields. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because inoculants should not replace soil testing or adequate fertility, and poor-quality or poorly matched products may show little field benefit, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Modern Tools and Future

DNA sequencing and metagenomics allow researchers to detect organisms that cannot be cultured easily. Metatranscriptomics, metabolomics and stable-isotope methods can help connect community composition with

function. Practical implementation of modern tools and future direction should start with a clear baseline and a defined management objective. For microbiome-aware sustainable crop and soil management, priority should be given to linking sequencing, metagenomics and functional measurements with practical agronomy rather than simply cataloguing organisms. A practical plan is to combine molecular data with soil chemistry, crop performance and management history, and validate proposed indicators across sites and seasons. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include community composition, functional genes, enzyme activity, nutrient cycling, disease outcomes and reproducibility. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that large datasets can reveal correlations that do not necessarily prove causation, so field experiments remain essential. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

The future goal is not simply to list microorganisms but to identify reliable microbiome indicators and management practices that improve crop performance across environments. Long-term field experiments are essential for this transition. The value of modern tools and future direction also depends on integration with the wider production system. Management should connect linking sequencing, metagenomics and functional measurements with practical agronomy rather than simply cataloguing organisms with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for community composition, functional genes, enzyme activity, nutrient cycling, disease outcomes and reproducibility.

Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because large datasets can reveal correlations that do not necessarily prove causation, so field experiments remain essential, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Conclusion

The soil microbiome is a living component of soil fertility. Microorganisms support nutrient cycling, root symbiosis, organic-matter turnover and disease suppression, and their activity is strongly influenced by crop and soil management.

Sustainable crop production can support beneficial microbial functions through diversity, organic inputs, balanced fertility and appropriate disturbance. Microbial products can add value, but they are most effective when integrated with good agronomy and locally tested soil management.

References

- Berendsen, R. L., Pieterse, C. M. J., & Bakker, P. A. H. M. (2012). The rhizosphere microbiome and plant health. *Trends in Plant Science*, 17, 478-486.
- Bulgarelli, D., Schlaeppi, K., Spaepen, S., van Themaat, E. V. L., & Schulze-Lefert, P. (2013). Structure and functions of the bacterial microbiota of plants. *Annual Review of Plant Biology*, 64, 807-838.
- Philippot, L., Raaijmakers, J. M., Lemanceau, P., & van der Putten, W. H. (2013). Going back to the roots: The microbial ecology of the rhizosphere. *Nature Reviews Microbiology*, 11, 789-799.
- Toju, H., Peay, K. G., Yamamichi, M., et al. (2018). Core microbiomes for sustainable agroecosystems. *Nature Plants*, 4, 247-257.
- Trivedi, P., Leach, J. E., Tringe, S. G., Sa, T., & Singh, B. K. (2020). Plant-microbiome interactions: From community assembly to plant health. *Nature Reviews Microbiology*, 18, 607-621.
- van der Heijden, M. G. A., Bardgett, R. D., & van Straalen, N. M. (2008). The unseen majority: Soil microbes as drivers of plant diversity and productivity. *Ecology Letters*, 11, 296-310.