

Regenerative Agriculture for Restoring Soil Health

Manjot Singh^{*1} and Shagun²

¹Ph.D., Soil Science; Lecturer, Surender Kaur Memorial Agriculture College, 24 BB, Padampur, Sri Ganganagar, Rajasthan, India

²M.Sc., Department of Vegetable Science, Dr. Yashwant Singh Parmar University of Horticulture and Forestry, Nauni, Solan, Himachal Pradesh, India

*Corresponding Author Email: manjotchinzer97@gmail.com

Abstract

Regenerative agriculture is a broad management approach that aims to rebuild soil function while maintaining productive farming. Although definitions vary, commonly emphasised practices include reducing unnecessary soil disturbance, keeping the soil covered, increasing crop diversity, maintaining living roots for more of the year, recycling organic materials and, where appropriate, integrating livestock. These practices can influence soil organic matter, aggregation, infiltration, erosion, nutrient cycling and biological activity. Soil restoration is not achieved by a single technique and outcomes depend on climate, soil type, crop system and starting condition. Reduced tillage may protect aggregates but can require changes in weed management. Cover crops add roots and residues but require water and careful termination. Organic amendments can increase carbon and nutrients but must be applied at agronomically appropriate rates. Regenerative systems should therefore be evaluated with measurable indicators such as soil organic carbon, aggregate stability, infiltration, bulk density, nutrient status, biological activity and crop yield stability. Economic performance and labour requirements are also important because practices must be maintained over many years to influence soil properties. Regenerative agriculture is most useful when treated as an adaptive process: farmers combine locally suitable practices, monitor results and modify the system rather than following a fixed recipe.

Keywords: Regenerative Agriculture, Soil Health, Cover Crops, Reduced Tillage, Soil Organic Matter, Crop Diversity

How to Cite this article: Singh, M. (2026). Regenerative Agriculture for Restoring Soil Health. *Future Agriculture e-magazine*, September 2026, Issue 3, pp. 303-310.

Introduction

Regenerative agriculture has gained attention as farmers seek production systems that improve rather than degrade soil resources. The term does not have one universal definition, but most approaches focus on rebuilding soil physical, chemical and biological function through changes in cropping and residue management. Practical implementation of introduction should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to combining reduced disturbance, cover, diversity, living roots and organic recycling into a locally adapted soil-restoration strategy. A practical plan is to begin

with clearly defined soil constraints and a baseline for erosion, organic matter, compaction, infiltration, fertility and crop performance. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include soil organic carbon, cover, bulk density, infiltration, aggregate stability, yield, input use and profitability. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that the term regenerative agriculture is broad, so claims should be tied to measurable changes rather than labels alone. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary,

technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Soil restoration is gradual. A farm with erosion, compaction or low organic matter may need several complementary practices and multiple seasons before measurable change occurs. The practical goal is to create a system that remains productive while improving resilience. The value of introduction also depends on integration with the wider production system. Management should connect combining reduced disturbance, cover, diversity, living roots and organic recycling into a locally adapted soil-restoration strategy with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for soil organic carbon, cover, bulk density, infiltration, aggregate stability, yield, input use and profitability. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because the term regenerative agriculture is broad, so claims should be tied to measurable changes rather than labels alone, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Minimising Unnecessary Disturbance

Frequent intensive tillage breaks aggregates, leaves soil exposed and can Practical implementation of minimising unnecessary disturbance should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to reducing aggressive tillage while maintaining seed placement, drainage, weed control and trafficability. A practical plan is to adjust tillage intensity gradually, use suitable seeding equipment, manage wheel traffic and monitor compaction rather than applying zero tillage as a universal rule.

Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include fuel use, residue cover, emergence, penetration resistance, bulk density, weeds, erosion and yield. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that reduced tillage can shift weed communities or create surface compaction in some soils if the rest of the system is not redesigned. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

accelerate organic-matter decomposition. Reduced tillage or no-till can lower disturbance and protect surface residues. The response depends on soil texture, drainage and machinery. The value of minimising unnecessary disturbance also depends on integration with the wider production system. Management should connect reducing aggressive tillage while maintaining seed placement, drainage, weed control and trafficability with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for fuel use, residue cover, emergence, penetration resistance, bulk density, weeds, erosion and yield. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because reduced tillage can shift weed communities or create surface compaction in some soils if the rest of the system is not redesigned, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Reduced tillage may increase weed-management challenges or surface compaction in some systems. Regenerative management therefore focuses on minimising unnecessary disturbance rather than treating zero tillage as a universal rule.

Keeping Soil Covered

Mulch and crop residues protect against raindrop impact, reduce evaporation and moderate soil temperature. Cover crops can provide living protection between cash crops and contribute additional root and shoot biomass. Practical implementation of keeping soil covered should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to using residues, mulch and cover crops to protect against erosion, evaporation and temperature extremes. A practical plan is to balance residue retention with livestock-feed needs, choose cover crops suited to rainfall and planting windows, and manage termination so the next crop is not disadvantaged. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include ground cover, soil moisture, surface temperature, erosion, cover biomass, establishment and subsequent crop yield. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that cover crops use water and nutrients as well as provide benefits, making species choice and termination timing important in dry environments. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

The amount of residue that can remain on the field depends on competing uses such as livestock feed. Partial retention or dedicated cover crops may be a practical compromise in mixed farming systems. The value of keeping soil covered also depends on integration with the wider production system. Management

should connect using residues, mulch and cover crops to protect against erosion, evaporation and temperature extremes with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for ground cover, soil moisture, surface temperature, erosion, cover biomass, establishment and subsequent crop yield. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because cover crops use water and nutrients as well as provide benefits, making species choice and termination timing important in dry environments, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Table 1. Common regenerative agriculture principles and soil-health effects.

Principle	Potential soil-health effect
Minimise disturbance	Protects aggregates and can reduce erosion and rapid organic-matter loss.
Maintain soil cover	Reduces erosion, evaporation and temperature extremes.
Increase crop diversity	Broadens root architecture, residue quality and biological interactions.
Maintain living roots	Provides continuous rhizosphere carbon and supports soil biology.
Recycle organic materials	Adds carbon and nutrients and supports aggregation.
Integrate livestock carefully	Recycles nutrients and diversifies production when grazing is well managed.

Crop Diversity and Living Roots

Different crops support different rooting patterns, nutrient demands and microbial communities. Rotations that include legumes, cereals and deep-rooted species can interrupt pest cycles and increase the diversity of organic inputs. Practical implementation of crop diversity and living roots should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to increasing rotation diversity and the duration of active roots to support soil structure, nutrient cycling and biological activity. A practical plan is to include cereals,


legumes and other functional crop types where practical, and use cover or relay crops to fill otherwise bare periods. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include rotation length, root biomass, soil cover days, nutrient use, pest pressure, microbial activity and yield stability. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that diversity should support agronomic and market objectives; adding species without a clear role can increase cost and management complexity. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Maintaining living roots for more of the year supplies carbon to rhizosphere microorganisms. Cover crops, relay crops and intercropping can extend root activity, but their water use and termination timing must be managed carefully in dry regions. The value of crop diversity and living roots also depends on integration with the wider production system. Management should connect increasing rotation diversity and the duration of active roots to support soil structure, nutrient cycling and biological activity with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for rotation length, root biomass, soil cover days, nutrient use, pest pressure, microbial activity and yield stability. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because diversity should support agronomic and market objectives; adding species without a clear role can increase cost and management complexity, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement

from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Organic Matter and Nutrient Cycling

Compost, manure and crop residues add carbon and nutrients. Organic matter improves aggregation and water-holding capacity and provides energy for soil organisms. However, nutrient release from organic materials may not match crop demand exactly. Practical implementation of organic matter and nutrient cycling should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to building organic matter through crop residues, compost, manure and root inputs while matching nutrients to crop demand. A practical plan is to analyse or estimate amendment nutrient content, integrate with soil tests and fertiliser plans, and avoid repeated over-application.



Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include organic carbon, nutrient balance, mineral nitrogen, phosphorus, aggregation, water holding and crop response. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that organic amendments can create nutrient loss, salinity, weeds or pathogen risks if quality and application rates are not controlled. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist.

Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Soil testing remains necessary. Manure and compost should be analysed or estimated so that phosphorus and nitrogen are not applied repeatedly above crop requirement. Regenerative systems still require nutrient budgeting. The value of organic matter and nutrient cycling also depends on integration with the wider production system. Management should connect building organic matter through crop residues, compost, manure and root inputs while matching nutrients to crop demand with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for organic carbon, nutrient balance, mineral nitrogen, phosphorus, aggregation, water holding and crop response. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because organic amendments can create nutrient loss, salinity, weeds or pathogen risks if quality and application rates are not controlled, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Integrating Livestock Where Appropriate

Managed grazing can recycle nutrients and convert cover-crop biomass into animal products. Short grazing periods followed by adequate recovery may fit some regenerative systems and can reduce the need for mechanical residue management. Practical implementation of integrating livestock where appropriate should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to using managed grazing to recycle nutrients and harvest cover-crop biomass without sacrificing

soil cover or structure. A practical plan is to control stocking density, grazing duration and recovery time, and avoid grazing when soil is too wet or plants are vulnerable. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include residual cover, compaction, infiltration, manure distribution, animal performance and regrowth. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that poorly timed or excessive grazing can reverse soil-health gains by exposing soil, compacting wet fields and reducing plant recovery. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Poorly managed grazing can compact wet soil, reduce cover and damage plants. Stocking density, duration and recovery period should therefore be adjusted to forage availability and soil condition. The value of integrating livestock where appropriate also depends on integration with the wider production system. Management should connect using managed grazing to recycle nutrients and harvest cover-crop biomass without sacrificing soil cover or structure with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for residual cover, compaction, infiltration, manure distribution, animal performance and regrowth. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because poorly timed or excessive grazing can reverse soil-health gains by exposing soil, compacting wet fields and reducing plant recovery, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal

variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Table 2. Practical indicators for tracking soil restoration.

Indicator	What it helps assess
Soil organic carbon	Long-term change in soil organic matter and carbon storage.
Aggregate stability	Resistance of soil structure to breakdown by water or disturbance.
Infiltration rate	Ability of soil to absorb rainfall or irrigation.
Bulk density	Compaction and pore-space condition.
Soil pH and nutrients	Chemical constraints and nutrient balance.
Yield and yield stability	Whether soil improvements translate into reliable farm production.

Water, Erosion and Soil Structure

Improved ground cover and aggregation can increase infiltration and reduce runoff. Deep and diverse root systems create channels that support water movement and can improve access to subsoil moisture. Practical implementation of water, erosion and soil structure should start with a clear baseline and a defined management objective.



For regenerative agriculture for measurable restoration of soil health, priority should be given to improving infiltration, aggregate stability and surface protection so rainfall is captured rather than lost as runoff. A practical plan is to maintain cover, reduce crusting and traffic damage, manage slopes and drainage, and use field observations or simple infiltration tests to identify weak areas. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include infiltration rate, runoff, erosion, aggregate stability, bulk density, soil moisture and yield during dry periods. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is

that better infiltration is valuable only when drainage and waterlogging risks are also considered, particularly in heavy soils. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Infiltration benefits are important but should be measured rather than assumed. Soil crusting, compaction layers and sodicity may require additional interventions. Contour farming, grassed waterways or terraces can complement regenerative field practices on erosion-prone land. The value of water, erosion and soil structure also depends on integration with the wider production system. Management should connect improving infiltration, aggregate stability and surface protection so rainfall is captured rather than lost as runoff with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for infiltration rate, runoff, erosion, aggregate stability, bulk density, soil moisture and yield during dry periods. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because better infiltration is valuable only when drainage and waterlogging risks are also considered, particularly in heavy soils, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Measuring Soil Health

Soil health should be assessed with a group of indicators. Organic carbon, bulk density, aggregate stability, infiltration, pH and nutrient status provide different information. Biological measures such as respiration or microbial biomass can add value when interpreted carefully. Practical implementation of measuring soil health should start with a clear

baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to tracking a practical group of physical, chemical and biological indicators rather than relying on a single soil-health score. A practical plan is to sample consistently, use repeatable field tests and link soil indicators with crop yield, rooting and management history. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include organic carbon, bulk density, aggregate stability, infiltration, pH, available nutrients, biological activity and yield stability. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that many indicators vary seasonally, so comparisons should use similar timing, depth and methods over multiple years. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Crop yield and yield stability remain important outcomes. A practice that improves one soil indicator but creates persistent yield or economic losses may not be sustainable for the farm. Repeated measurements over time are more useful than a single test. The value of measuring soil health also depends on integration with the wider production system. Management should connect tracking a practical group of physical, chemical and biological indicators rather than relying on a single soil-health score with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for organic carbon, bulk density, aggregate stability, infiltration, pH, available nutrients, biological activity and yield stability. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because many

indicators vary seasonally, so comparisons should use similar timing, depth and methods over multiple years, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Economics and Adaptive Management

Many regenerative practices involve transition costs, learning and changes in equipment or labour. Cover-crop seed and additional management may increase short-term cost, while benefits such as erosion control and improved water retention may accumulate over time. Practical implementation of economics and adaptive management should start with a clear baseline and a defined management objective. For regenerative agriculture for measurable restoration of soil health, priority should be given to ensuring that soil restoration can be maintained through realistic labour, machinery, seed and input costs. A practical plan is to start with priority fields, compare strips or management zones, record transition costs and adjust practices according to measured agronomic and financial outcomes. Changes should be introduced consistently so their effects can be interpreted. Workers should know the normal range and the signs that require attention. Monitoring can include gross margin, input cost, labour, machinery, yield stability, risk exposure and soil-health indicators. Trends are more useful than isolated readings because season, production stage and local conditions alter expected performance. One limitation is that benefits may take several seasons to emerge, so short-term economics and long-term resource improvement should be considered together. Therefore records should be reviewed regularly and important deviations checked by the relevant farm, veterinary, technical or crop specialist. Measures that fit available labour, infrastructure and budget are more likely to be maintained and to produce reliable long-term benefit.

Farmers should start with clearly defined objectives and compare treated and untreated areas where practical. Adaptive management

allows practices to be modified when weather, markets or soil response differ from expectations. The value of economics and adaptive management also depends on integration with the wider production system. Management should connect ensuring that soil restoration can be maintained through realistic labour, machinery, seed and input costs with routine records, clear responsibilities and a response plan. Where equipment, biological processes or field operations are involved, timing, maintenance and quality control matter as much as the initial choice. Records should show what was done and the response observed, particularly for gross margin, input cost, labour, machinery, yield stability, risk exposure and soil-health indicators. Evaluation should consider productivity together with health or soil function, labour, input use, risk and economics. Because benefits may take several seasons to emerge, so short-term economics and long-term resource improvement should be considered together, weak results should first prompt a review of implementation and local constraints. Comparison with a previous season, baseline or similar group helps separate real improvement from normal variation. This adaptive process allows timing, thresholds and resource use to be refined as experience accumulates.

Conclusion

Regenerative agriculture provides a framework for restoring soil health through reduced disturbance, soil cover, diversity, living roots and organic matter management. Its success depends on how these principles are adapted to local soil, climate and farming goals.

The most credible regenerative systems use measurable soil and crop indicators and recognise trade-offs. Long-term improvement comes from consistent management and learning rather than from any single practice or product.

References

- Giller, K. E., Hijbeek, R., Andersson, J. A., & Sumberg, J. (2021). Regenerative agriculture: An agronomic perspective. *Outlook on Agriculture*, 50, 13-25.
- Lal, R. (2020). Regenerative agriculture for food and climate. *Journal of Soil and Water Conservation*, 75, 123A-124A.
- Newton, P., Civita, N., Frankel-Goldwater, L., Bartel, K., & Johns, C. (2020). What is regenerative agriculture?

- A review of scholar and practitioner definitions. *Frontiers in Sustainable Food Systems*, 4, 577723.
- Schreefel, L., Schulte, R. P. O., de Boer, I. J. M., Schrijver, A. P., & van Zanten, H. H. E. (2020). Regenerative agriculture: The soil is the base. *Global Food Security*, 26, 100404.
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G. P., & Smith, P. (2016). Climate-smart soils. *Nature*, 532, 49-57.
- Pittelkow, C. M., Liang, X., Linquist, B. A., et al. (2015). Productivity limits and potentials of conservation agriculture. *Nature*, 517, 365-368.